

This product fulfills the sanitizing criteria of the US PUBLIC HEALTH SERVICE in waters up to 500 ppm of hardness calculated as CaCO₃ when tested by the AOAC Germicidal and Detergent Sanitizers - Official Method.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS and DOMESTIC ANIMALS

WARNING: Tablet dust causes substantial but temporary eye injury. Do not get dust in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Tablets may be harmful if swallowed.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN: Rinse immediately with water for 15-20 minutes. If irritation persists, call poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by poison control center doctor. Do not give anything by mouth to an unconscious person. Have the product label or container with you when calling a poison control center or doctor, or going for treatment.

STORAGE and DISPOSAL: Do not contaminate water, food or feed by storage and disposal. Store in original container in areas inaccessible to persons unfamiliar with its use. Nonrefillable container. Do not reuse or refill this container. Triple rinse promptly after emptying. Offer empty container for recycling if available or discard in trash. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

STERAMINE-0113

STERAMINE™

www.sanitize.com

6-Q Tablets

The Multi-Purpose Sanitizer



FOR SANITIZING FOOD CONTACT SURFACES

USE ONE TABLET PER 1½ GALLONS OF WATER

Kills HIV-1 (AIDS Virus) when used as directed for sanitizing.

EPA REG. NO. 1561-12

Active ingredient (Quaternary)	50%
Alkyl (C14 95%, C12 3%, C16 2%) dimethyl benzyl ammonium chloride dihydrate	50%
INERT INGREDIENTS	50%
TOTAL	100%

WARNING

KEEP OUT OF REACH OF CHILDREN

See left panel for additional precautionary statements. NET WEIGHT 8 OUNCES
For Institutional and Commercial use only.

100
Tablets

Manufacturing Chemists
EDWARDS-COUNCILOR CO. INC., 1427 Baker Road, Virginia Beach, VA 23455

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Prepare sanitizing solutions with warm water. Allow several minutes for tablets to dissolve before using.

FOR SANITIZING DISHES, GLASSES, AND UTENSILS IN RESTAURANTS, TAVERNS, ETC.

1. Scrape and prewash utensils and glasses whenever possible.
2. Wash with a good detergent or compatible cleaner in first sink compartment.
3. Rinse with clean water in second sink compartment.
4. Sanitize in a solution of 1 TABLET per 1½ GALLONS OF WATER (200 ppm) in third sink compartment. Immerse all utensils for at least one minute or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drainboard to air dry.
6. A fresh sanitizing solution must be prepared at least daily or more often if the solution becomes diluted or soiled.

DIRECTIONS FOR SPRAYING

FOR SANITIZING FOOD PROCESSING EQUIPMENT, DAIRY EQUIPMENT, SINKS, COUNTERTOPS, REFRIGERATED STORAGE AND DISPLAY EQUIPMENT and other hard non-porous food contact articles and surfaces.

1. Wash and rinse all articles and surfaces thoroughly.
2. Apply a solution (SPRAY ON WITH HAND TRIGGER SPRAYER OR WIPE ON) of 200 ppm concentration and allow surface to remain wet for at least one minute followed by adequate draining and air drying.

KILLS HIV-1 (AIDS Virus) when used as directed for sanitizing.

EPA-Est. 1561-VA-1

